

Available online at www.sciencedirect.com

ScienceDirect

Journal of Sport and Health Science xx (2016) 1–8

www.jshs.org.cn

Original Article

Intention and automaticity toward physical and sedentary screen-based leisure activities in adolescents: A profile perspective

Gonzalo Marchant ^{a,b,†}, Guillaume Chevance ^{a,c,†}, Julie Boiché ^{a,*}^a *Laboratory Epsilon “Dynamics of Human Abilities and Health Behavior” (EA 4556), University of Montpellier, 4 Boulevard Henri IV, Montpellier 34000, France*^b *Laboratory L-ViS (EA 7428), Université Claude Bernard Lyon 1, Campus de La Doua, Bâtiment R. Dubois - 27–29 bd du 11 novembre 1918, Villeurbanne cedex 69622, France*^c *Clinique du Souffle LaVallonie, 5 Santé Group, 800, Avenue Joseph Vallot, Lodève 34700, France*

Received 4 February 2015; revised 16 July 2015; accepted 19 April 2016

Available online

Purpose: Physical activity (PA) and sedentary behavior (SB) are increasingly considered independent health behaviors. Additionally, current research suggests that both controlled and automatic determinants account for their adoption. The purpose of this article is to identify intention–automaticity profiles toward PA and screen-based SB and to examine how those profiles are associated with different behavioral patterns.

Method: Two cross-sectional studies based on self-report questionnaires were conducted with French high school students (Study 1: $n = 198$; Study 2: $n = 185$).

Results: In all, 4 distinct motivational profiles appeared. The first 3 clusters emerged in both studies: “PA” (high levels of automaticity and intention for PA, low levels of automaticity and intention for screen-based SB); “screen” (high levels of automaticity and intention for screen-based SB, low levels of automaticity and intention for PA), and “mixed” (high levels of all variables), whereas the fourth cluster was observed only in Study 2: “high control” (below-mean levels of automaticity, high levels of intention toward both PA and screen-based SB). Adolescents with a screen profile displayed the least healthy behavioral pattern, whereas those in the PA profile demonstrated the most favorable behaviors.

Conclusion: Future research is needed to extend these results to other populations using complementary assessment methods of automatic psychological processes and PA and SB behaviors.

© 2016 Production and hosting by Elsevier B.V. on behalf of Shanghai University of Sport. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Active lifestyle; Cluster analysis; Exercise; HabitMotivation; Physical activity

1. Introduction

Increasing evidence indicates that physical activity (PA) levels in youth have declined during the past decades, with currently 4 of 5 adolescents not reaching the levels recommended in public health guidelines.¹ Meanwhile, past research also indicates that the time devoted to sedentary behavior (SB) has gradually increased over time, particularly in children and adolescents.² Whereas for a long time PA and SB have been regarded as functional opposites—for example, through the perspective of the “displacement hypothesis”³—they tend more and more to be conceived as 2 distinct categories of behaviors, which are likely to have independent effects on health.⁴ In this

vein, the mean durations devoted to those behaviors were previously found to be mainly independent and only marginally negatively related,⁵ in particular in the case of screen-based SB and leisure-time PA. This suggests that various patterns of behaviors could exist in the population. In particular, young people could combine high levels of both PA and SB. In this regard, several authors suggest that those behaviors present at least partly distinct psychological determinants.⁶

Recently, it was advocated by health psychologists that behavior adoption relies on 2 main kinds of processes, that is, controlled processes, which imply deliberate decisions and intentions, and automatic processes, characterized by “4 horse-men”, namely, lack of awareness, efficiency, lack of intention, and difficulty in controlling one’s behavior.⁷ Automaticity is not conceived as an “all-or-nothing” characteristic, but psychological processes can be considered to lie on a continuum. More particularly, each process can be deemed to occur with more or less awareness on the part of the individual, to be more or less

Peer review under responsibility of Shanghai University of Sport.

* Corresponding author.

E-mail address: julie.boiche@umontpellier.fr (J. Boiché).

† The first two authors equally contributed to the article.

<http://dx.doi.org/10.1016/j.jshs.2016.08.006>

2095-2546/© 2016 Production and hosting by Elsevier B.V. on behalf of Shanghai University of Sport. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

easy to control, to be more or less efficient, and to be (un)intentional, to some degree.⁸ In previous research conducted on habits, it was advanced that individuals are unlikely to be aware of some of the processes of engaging in a health-related behavior,⁹ even when the behavior is characterized as being fully conscious.¹⁰ However, meta-cognitive instruments can be used based on the assumption that people are able to reflect the other features of automaticity—that is, lack of intention, control, and attention, particularly for exercise.¹¹

Among controlled processes, intentions have been established as the strongest construct associated with PA in adolescents.¹² In the past decades, several studies investigated this relationship, in particular based on the theory of planned behavior.¹³ In 2002, the results of 72 studies were pooled in a meta-analysis, which supported the tenets of the theory of planned behavior in the case of PA (i.e., intentions mediate the relationships between attitudes, social norms, perceived behavioral control, and behavior adoption; perceived behavioral control also has a direct relationship with behavior).¹⁴ A more recent meta-analysis confirms that intentions are significantly associated with PA behavior, controlling for perceived behavioral control and past PA behavior.¹⁵ Even if SB were less studied, a handful of studies also indicated that intentions to adopt various SBs, including TV watching, are significantly associated with the time spent doing them.¹⁶ Recently, health psychologists were compelled to simultaneously consider controlled and automatic processes.¹⁷ Indeed, behaviors need to be repeated to have significant effects on health, which is particularly relevant in the case of PA. In other words, health-related behaviors are deemed more beneficial when they become habits. Whereas assessing the frequency of behavior was long considered the best strategy to capture habits, a more contemporary view proposes to consider this construct as a repeated behavior that has gained a degree of automaticity and is executed in stable contexts.¹¹

Numerous investigations of the role played by the automatic properties involved in habitual behavior were conducted in the past decade, thanks to the development of self-report scales to capture automatic processes, in particular the Self-Report Habit Index (SRHI).¹⁸ A recent meta-analysis examined the links between habit strength and health behavior adoption, including PA and SB.¹⁹ It revealed the existence of medium to strong correlations between habit strength and behavior adoption. It also supported the theory that the contribution of intentions and automaticity levels in the explanation of PA and SB operated in interaction. More precisely, intentions tend to show lower associations with behavior when habit levels are high, and vice versa. Furthermore, past research suggested that automaticity toward 1 behavior may affect not only the frequency of its adoption but also individuals' engagement in other behaviors.²⁰ More precisely, it was observed that a strong habit toward television viewing was positively associated with soft drink consumption. Automatic processes could thus be seen as potential predictors of a pattern of interconnected behaviors rather than the determinants of a single, separated action.

In summary, it seems that both intentional and automatic psychological processes are worthy of consideration when

trying to understand the adoption of health-related behaviors such as PA and SB. Additionally, there is increasing evidence that those categories of behavior are partly independent, but it was also advanced that they could share common determinants. In particular, it is possible that the psychological dispositions toward 1 kind of behavior could affect a different category of behavior. In other words, cross-context correlations should be expected (i.e., between motivational determinants of PA with SB, and vice versa). Considering those complex assumptions, the current research adopted a person-centered approach. More specifically, the purposes of the present study were (1) to identify intention–automaticity profiles toward PA and screen-based SB in adolescents and (2) to examine how those profiles were associated with the adoption of significantly different behavioral patterns in terms of PA and screen-based SB.

2. Study 1

2.1. Methods

2.1.1. Participants and procedure

A cross-sectional design was conducted for 198 10th grade students (108 females, 70 males, and 20 students who did not provide gender) from a general French high school (age range 14–20 years; 16.00 ± 0.76 , mean \pm SD). The study protocol was in accordance with the principles of the Declaration of Helsinki of 1975, as revised in 2000. It was approved by the Ethics Committee research team at the University of Montpellier, France. All participants signed the informed consent presenting the purpose of the study, which indicated that their participation was voluntary and that they could cease it at any time. They were assured that confidentiality would prevail during the whole study and that only the means of the participants would be analyzed and communicated. Questionnaire completion occurred during a compulsory physical education lesson supervised by a teacher and lasted about 15 min.

2.1.2. Measures

2.1.2.1. PA. The International Physical Activity Questionnaire was used.²¹ The participants were asked to indicate (1) how many days they engaged in at least 10 min of vigorous PA, moderate PA, and walking during the previous week, and (2) the average duration of those sessions. The International Physical Activity Questionnaire has acceptable measurement properties for monitoring levels of habitual PA in male and female adolescents older than 14 years.²²

2.1.2.2. Screen-based SB. The time spent by participants watching TV or using a computer was assessed by means of the corresponding 2 items of the Sedentary Behavior Questionnaire,²³ distinct questions having been formulated regarding weekdays and weekends. The total amount of time per week was computed by multiplying the number of days in which they adopted such behaviors by the average duration of those sessions. The 2-week reliability and validity of those items was demonstrated in adults,²³ and they were successfully used in previous research among high school students.²⁴

2.1.2.3. *Automaticity.* Automaticity toward PA and SB was measured with the Self-Report Behavioral Automaticity Index (SRBAI).²⁵ Following the stem “Behavior *X* is something. . .,” this 4-item scale (“that I do without thinking”; “that I do automatically”; “that I do without having to consciously remember”; “that I start doing before realizing I am doing it”) includes a Likert scale ranging from 1 (*do not agree at all*) to 5 (*totally agree*). Internal consistency was high for both PA ($\alpha = 0.80$) and screen-based SB ($\alpha = 0.80$).

2.1.2.4. *Intentions.* Intentions were measured with 2 items (“Do you have the intention to practice regular physical activity, watch TV, or use a computer regularly during the upcoming week?”; “Are you sure to practice regular physical activity, watch TV, or use a computer regularly during the upcoming week?”). Participants answered on a 5-point bipolar scale ranging from -2 (no, not at all) to $+2$ (yes, absolutely). Internal consistency was high for both behaviors (PA, $\alpha = 0.85$; screen-based SB, $\alpha = 0.90$).

2.1.3. Data analysis

Regarding items for which less than 5% of data were missing, a replacement by the mean score was applied ($n = 45$).²⁶ Complete data were available for 147 participants. To reach the study’s first objective, a series of cluster analyses were conducted to create groups of participants according to their intention and automaticity profiles. To avoid distortion in the formation of clusters, particular attention should be paid to metrics, outliers, and multicollinearity.²⁷ After these prerequisites were verified, a hierarchical cluster analysis was performed to explore the number of clusters that naturally emerge, using Ward’s method as algorithm and squared Euclidean distance as similarity measure. The obtained centroids’ characteristics were then used in a confirmatory k-mean cluster analysis with a defined number of clusters. Next, ANOVAs were conducted to test the intercluster differences in terms of automaticity and intention. Last, a series of ANOVAs was performed to observe potential differences between clusters in terms of behavior adoption.

2.2. Results

2.2.1. Descriptive statistics

Correlations between the variables appear in Table 1. A correlation for the data revealed that intentions and automaticity scores were significantly but not highly related ($r = 0.56$, $r = 0.42$ for PA and screen-based SB, respectively, $p < 0.05$). PA intentions and automaticity were significantly associated with high and moderate intensity PA ($0.27 < r_s < 0.56$, $p_s < 0.05$) but not with walking ($r = 0.05$, $r = 0.10$, respectively, $p > 0.05$). Screen-based SB intentions and automaticity were significantly correlated to the time spent in front of a screen ($r = 0.29$, $r = 0.44$, respectively, $p < 0.05$). Last, cross-context correlations (i.e., between motivational determinants of PA and screen-based SB, or vice versa) were nonsignificant or low. For instance, the higher the level of PA automaticity in participants, the lower their intentions to spend time in front of a television or a computer ($r = -0.27$, $p < 0.05$).

2.2.2. Prerequisites to cluster analysis

The mean \pm SD values for each variable of the study appear in Table 2. The variables included in the cluster analysis were standardized so as to contribute equally to the formation of the clusters.²⁷ Five univariate outliers were excluded from further analysis (distance from the mean superior to 3.29 SD units), as well as 6 multivariate outliers (Mahalanobis distance significant at the $p = 0.001$ threshold). Finally, because the highest correlation was equal to $r = 0.51$, no problem of multicollinearity was encountered.

2.2.3. Cluster analysis

A hierarchical cluster analysis was performed entering intention and automaticity toward PA and SB. A large increase of the agglomeration coefficient (29%) suggested a 3-cluster solution to be suitable. This analysis was followed by a confirmatory k-mean cluster analysis. The centroids’ characteristics appear in Table 2, and Fig. 1A describes the 3 intention–automaticity profiles. The first cluster was labeled “screen”, because students in this group were characterized by high levels of automaticity and intention for screen-based SB, but low

Table 1
Correlation coefficients for Studies 1–2.

	1	2	3	4	5	6	7	8
Physical activity								
1. Intention	—	0.54*	0.57*	0.28*	0.01	-0.22*	-0.28*	-0.40*
2. Automaticity/lack of control	0.56*	—	0.40*	0.22*	0.00	-0.31*	-0.07	-0.33*
3. High intensity	0.48*	0.49*	—	0.12	-0.08	-0.27*	-0.12	-0.24
4. Moderate intensity	0.33*	0.27*	0.24*	—	0.23*	-0.01	-0.05	-0.09
5. Walking	0.05	0.10	0.16	-0.07	—	-0.04	0.06	0.10
Sedentary behavior								
6. Intention	-0.17	-0.27*	-0.01	0.02	0.09	—	0.29*	0.51*
7. Automaticity/lack of control	0.06	-0.02	-0.10	0.01	-0.01	0.42*	—	0.37*
8. Screen-based time	0.15	-0.03	0.16	0.14	0.05	0.29*	0.44*	—

Note: Correlation coefficients for Study 1 are selected in bold.

* Correlation is significant at the $p < 0.05$ level.

Table 2
Mean value and centroid characteristics of intention and automaticity for PA and SB (Study 1, mean \pm SD).

	All <i>n</i> = 136		Mixed <i>n</i> = 49 (36%)		Screen <i>n</i> = 54 (40%)		Physical activity <i>n</i> = 33 (24%)		<i>F</i>	η^2	<i>p</i>
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>			
PA											
Intention	0.86 \pm 1.10		1.71 \pm 0.48		0.07 \pm 0.95*		1.06 \pm 0.77*#		59.39	0.47	0.000
Automaticity	2.95 \pm 1.08		3.65 \pm 0.80		2.04 \pm 0.68		3.30 \pm 0.76*#		66.07	0.50	0.000
High intensity	192.81 \pm 199.58		343.51 \pm 229.23		78.61 \pm 103.99*		213.20 \pm 142.96*#		26.85	0.34	0.000
Moderate intensity	93.65 \pm 159.87		127.67 \pm 203.46		36.07 \pm 55.03*		102.30 \pm 97.16*		4.79	0.08	0.010
Walking	203.23 \pm 243.47		236.01 \pm 303.46		140.97 \pm 113.04		212.50 \pm 193.60		2.16	0.04	0.120
SB											
Intention	1.18 \pm 0.98		1.62 \pm 0.53		1.65 \pm 0.42		0.63 \pm 0.67*#		44.99	0.40	0.000
Automaticity/LOC	3.05 \pm 1.10		3.67 \pm 0.84		3.36 \pm 0.97		1.98 \pm 0.46*#		44.89	0.40	0.000
Screen-based time	505.30 \pm 403.97		666.34 \pm 454.78		534.75 \pm 379.07		383.89 \pm 271.53*#		4.35	0.08	0.015

Note: *n* = 136, only 136 participants provided complete data on their behaviors.

* *p* < 0.05, compared with mixed group;

p < 0.05, compared with screen group.

Abbreviations: LOC = lack of control; PA = physical activity; SB = sedentary behavior.

levels of automaticity and intention for PA. It comprised 40% of the sample (*n* = 54). The second cluster was labeled “PA” and was characterized by high levels of automaticity and intention for PA, but low levels of automaticity and intention for

screen-based SB. It included 24% of the sample (*n* = 33). The third cluster, characterized by high levels of all variables, was labeled “mixed” and comprised 36% of the sample (*n* = 49).

A series of ANOVAs indicated significant differences among clusters regarding all psychological variables (Table 2). Newman-Keuls *post hoc* analyses indicated that the 3 groups were significantly distinct from each other on PA intentions (*p* < 0.05). However, regarding automaticity toward PA and intention and automaticity toward screen-based SB only the “physical activity” group obtained significantly different scores compared to the other two profiles.

2.2.4. Clusters characteristics

A series of ANOVAs indicated significant between-cluster differences regarding PA and screen-based SB. Participants in the mixed profile displayed significantly higher levels of vigorous PA, followed by those with the PA and the screen cluster. Participants with a screen profile also reported significantly lower levels of moderate-intensity PA compared with the other profiles. No statistical difference emerged regarding walking. Finally, participants in the mixed cluster reported the highest durations of screen-based SB, whereas those in the PA cluster showed the lowest scores.

3. Study 2

3.1. Method

Study 2 concerned 185 students (59 females; 126 males) recruited from grades 9 to 12 of a general French high school (age range 14–19 years; 17.00 \pm 0.83, mean \pm SD). The protocol, method, and data analysis strategies were similar to Study 1, with the exception of the measurement of automaticity. Three items assessing lack of control (LOC) were derived from the SRHI.¹⁸ A discriminant validity study indicates that the 4 items of the SRBAI fail to capture this facet of automaticity, contrary to those 3 SRHI items.²⁸ Following the stem “*Behavior X is something I do. . .*,” participants used a 5-point Likert scale ranging from 1 (*totally not agree*) to 5 (*totally agree*) to




Fig. 1. Intention-automaticity profiles (Study 1, A) and intention-lack of control profiles (Study 2, B) observed among secondary school students. HCR = high control profile; mixed = mixed profile; LOC = lack of control; PA = physical activity; SB = sedentary behavior; screen = screen profile.

characterize 3 affirmations (“that makes me feel weird if I do not do it”; “that would require effort not do it”; “that I would find hard not to do”). Internal consistency was satisfactory for all scales ($0.72 < \alpha < 0.93$). Regarding items for which less than 5% of data was missing, a replacement by the mean score was applied ($n = 23$).²⁶

3.2. Results

3.2.1. Descriptive statistics

Correlations between the variables appear in Table 1. There was a significant correlation among PA intention, PA LOC, and high intensity of PA. A significant, but small, negative association between PA intention and SB intention was observed. A small negative association between PA intention and LOC and screen-based time was also observed.

3.2.2. Prerequisites to cluster analysis

The mean \pm SD values for each variable appear in Table 3. The variables were standardized so as to contribute equally to the formation of the clusters.²⁷ No univariate outliers (distance from the mean superior to 3.29 SD units) were observed, but 1 multivariate outlier (Mahalanobis distance significant at the $p = 0.001$ threshold) was excluded. No problems of multicollinearity were encountered (highest correlation: $r = 0.51$).

3.2.3. Cluster analysis

A hierarchical cluster analysis was performed entering intention and LOC toward PA and screen-based SB. A large increase of the agglomeration coefficient (26%) suggested that a 4-cluster solution was suitable. A confirmatory k-mean cluster analysis was run. The centroids’ characteristics appear in Table 3, and Fig. 1B describes the 4 intention–LOC profiles.

Three clusters presented similar characteristics to those observed in study 1, namely a PA profile ($n = 65$, 34%), a screen profile ($n = 37$, 20%), and a mixed profile ($n = 36$, 20%). The fourth cluster was labeled high control because participants in this group displayed below-mean levels of LOC, but high levels

of intentions toward both PA and screen-based SB ($n = 46$, 26%).

A series of ANOVAs indicated significant differences between clusters for all psychological variables (Table 3). Newman-Keuls *post hoc* analyses indicated that the 4 groups were significantly distinct from each other on LOC toward PA, except between the PA and mixed profiles, and on LOC toward screen-based SB, except between the PA and the high control profiles. Regarding intention toward PA, the screen profile did not differ significantly from the other groups. Last, regarding intention toward screen-based SB, only the PA profile was significantly different from the others.

3.2.4. Cluster characteristics

A series of ANOVAs indicated significant between-cluster differences regarding PA and screen-based SB (Table 3). Newman-Keuls *post hoc* analyses indicated that participants in the screen group reported significantly lower levels of vigorous PA compared with the other profiles. No significant difference emerged regarding moderate PA or walking. Participants in the PA profile showed significantly lower levels of screen-based SB compared with the other groups.

4. Discussion

The purpose of this article was to identify intention–automaticity profiles with regard to PA and screen-based SB and to examine whether those profiles were associated with significantly different behavioral patterns in French adolescents. Study 1 underlined the presence of 3 motivational profiles: a screen and a PA cluster, characterized by high levels of both intentions and automaticity for 1 behavior (SB/PA) and low levels of those variables for the alternative (PA/SB), and a mixed cluster with high levels of both controlled and automatic forms of motivations for both behaviors. Those profiles were also observed in study 2, as well as a high control cluster characterized by high levels of intentions but low levels of LOC. Those results highlight the interest in studying the

Table 3
Mean value and centroid characteristics of intention and automaticity for PA and SB (Study 2).

	All $n = 184$		PA $n = 65$ (34%)		High control $n = 46$ (26%)		Mixed $n = 36$ (20%)		Screen $n = 37$ (20%)		F	η^2	p
	M	SD	M	SD	M	SD	M	SD	M	SD			
PA													
Intention	0.395 \pm 1.24		4.62 \pm 0.59		4.45 \pm 0.65*		4.12 \pm 0.92*#		1.96 \pm 0.80		119.08	0.66	0.000
LOC	2.89 \pm 1.18		3.54 \pm 0.51		2.60 \pm 1.00*		3.47 \pm 0.83#		1.53 \pm 0.68*#†		47.93	0.44	0.000
High intensity	289.24 \pm 289.25		411.10 \pm 320.89		298.48 \pm 250.96		315.61 \pm 277.27		60.67 \pm 104.46*#†		13.84	0.19	0.000
Moderate intensity	118.26 \pm 195.89		139.31 \pm 197.14		140.37 \pm 252.76		122.97 \pm 182.59		45.90 \pm 56.20		1.96	0.03	0.122
Walking	285.67 \pm 360.24		331.12 \pm 424.23		259.88 \pm 340.97		229.50 \pm 216.27 ^a		290.63 \pm 371.62		0.62	0.01	0.601
SB													
Intention	3.70 \pm 1.19		2.48 \pm 0.78		4.46 \pm 0.56*		4.43 \pm 0.75*		4.13 \pm 1.01*		82.39	0.58	0.000
/LOC	1.97 \pm 1.05		1.42 \pm 0.51		1.41 \pm 0.47		3.42 \pm 0.55*		2.25 \pm 1.26*#†		73.81	0.55	0.000
Screen-based time	789.20 \pm 77.03		375.35 \pm 232.34		832.67 \pm 695.64		1057.83 \pm 832.09		1267.33 \pm 1059.005		13.10	0.20	0.000

Note: $n = 184$, only 184 participants provided data on their behaviors.

* $p < 0.05$, compared with PA group;

$p < 0.05$, compared with high control group;

† $p < 0.05$, compared with mixed group.

Abbreviations: LOC = lack of control; PA = physical activity; SB = sedentary behavior.

1447 components of automaticity in health psychology, in particular
1448 pertaining to PA and SB. Indeed, recent work indicates that the
1449 4-item SRBAI mainly reflects the unintentional initiation and
1450 efficient realization of behavior, whereas the 3 items of the
1451 original SRHI retained in study 2 mostly reflect the LOC facet.
1452 The results also highlight the great consistency that appears in
1453 most adolescents between intentional and nonintentional pro-
1454 cesses. However, the correlation values between intention and
1455 automaticity scores suggest that automaticity is not all-or-
1456 nothing, but rather that intentions and automaticity can coexist.
1457 In other words, the results indicate that automaticity may not be
1458 a firm dichotomy, suggesting the existence of a continuum
1459 where the attributes of automaticity can develop gradually.
1460 Thus, if behavior initially was intentional and if intentions
1461 remain stable over time, then automaticity would correlate with
1462 intentions. In this vein, with the exception of the high control
1463 group in study 2, adolescents apparently tend to form intentions
1464 to adopt behaviors for which they also report strong levels of
1465 automaticity.

1466 The present work further provides evidence relative to the
1467 differentiated patterns of behaviors observed depending on ado-
1468 lescents' motivational profiles. Based on the observed correlations,
1469 the current set of studies confirms that both intentions and
1470 automaticity levels toward 1 type of behavior are strongly as-
1471 sociated with its adoption. Additionally, small correlations were
1472 obtained between PA and SB scores. On the one hand, the
1473 direction of these associations is consistent with the displace-
1474 ment hypothesis³; on the other hand, their small magnitude
1475 does not support the existence of a full substitution phenom-
1476 enon, highlighting the interest in studying both behaviors
1477 independently.⁵ Indeed, this result suggests that some individu-
1478 als can be at the same time sedentary and physically inactive
1479 or can present with simultaneously high physical activity and
1480 high sedentary time.²⁹ There were also several significant and
1481 negative cross-correlations between psychological processes
1482 (intention or automaticity/LOC) toward 1 behavior and the
1483 adoption of the other behavior. Thus, PA and screen-based SB
1484 share to a certain extent some common motivational determinants.

1485 Regarding the results relative to the association between
1486 motivational profiles and behaviors, Studies 1 and 2 revealed
1487 only partly similar patterns for PA. Indeed, whereas in Study 1
1488 the mixed profile was attached to higher levels of intense and
1489 moderate PA, in Study 2, students in the PA profile appeared
1490 to be the most active. Both studies indicated that the screen profile
1491 demonstrated the lowest levels of PA. It is noteworthy that no
1492 significant differences emerged for walking, suggesting that
1493 motivational profile mostly accounts for more intense PA.
1494 Regarding SB, the results consistently showed that adolescents
1495 with a screen profile spent the greatest amounts of time in this
1496 behavior, whereas those with a PA profile spent the least. Dis-
1497 playing a mixed or high control profile was associated with
1498 intermediate levels of SB. Ultimately, a PA profile appears to be
1499 associated with the healthiest behavioral pattern. On the other
1500 hand, participants with a mixed profile were found to be active
1501 but at the same time sedentary. These findings extend argu-
1502 ments that both types of motivation can be useful for explaining
1503 health behaviors, especially when automatic and controlled

1504 processes are expected to have separate mechanisms of behav-
1505 ioral regulation. These systems are inevitably intertwined as
1506 they regulate behavior.

1507 The results were not exactly similar in both studies regard-
1508 ing correlations, profiles, or association between profiles and
1509 behaviors. Several explanations may be advanced to account
1510 for those slightly different results. First, Study 1 and Study 2
1511 differed in terms of participants, with Study 2 including a
1512 wider range of grades. Because PA participation was previ-
1513 ously found to decrease and sedentary time to increase with
1514 age in adolescents,¹ this could account for varying results.
1515 Second, different items were used to evaluate automaticity, and
1516 Study 1 and Study 2 focused on different features of this
1517 construct.³⁰ This could explain why standardized scores were
1518 slightly different for the mixed profiles between both studies
1519 and why a fourth profile emerged.

1520 The current research supports the interest in a joint assess-
1521 ment of intention and automaticity for both PA and SB in
1522 adolescents. In line with recent research,⁵ it appears that various
1523 combinations of behaviors can be observed in youth, and
1524 knowing that an adolescent has adopted 1 behavior does not
1525 enable the reaching of conclusions on his or her involvement in
1526 the other. The present findings are, however, limited by several
1527 factors. A first limitation relies in the cross-sectional design. It
1528 would be interesting to conduct prospective studies to investi-
1529 gate whether intention–automaticity profile explains the PA and
1530 SB behaviors subsequently adopted. Another limitation is
1531 linked to the use of self-reported measurement, regarding both
1532 automaticity and behavior. Regarding automatic processes,
1533 recent debates occurred in the health psychology community
1534 regarding the nature of the processes actually captured by scales
1535 such as the SRHI or SRBAI.^{31,32} Indeed, this method implies
1536 meta-cognitive processes that cannot be considered purely auto-
1537 matic, and, in particular, it was advocated that individuals could
1538 not be expected to be able to recall phenomena occurring
1539 outside their consciousness. Some authors thus recommended
1540 the use of implicit tests, such as the implicit-association test or
1541 a “go–no go” task, to capture automatic processes, in the sense
1542 of their being out of one's awareness.³³

1543 Regarding PA and SB, individuals are likely to overestimate
1544 the time that they spend being active or, to the contrary, to
1545 underestimate the time that they spend watching TV or using
1546 a computer. In this regard, the use of objective assessment
1547 devices represents a fruitful research perspective. Indeed,
1548 devices such as accelerometers are increasingly being used to
1549 help young people estimate their actual energy expenditure.³⁴
1550 Intentional processes are also likely to predict structured activi-
1551 ties, whereas automatic processes would account to a greater
1552 extent for spontaneous behaviors.³⁵ Such a hypothesis could be
1553 explored in studies combining explicit self-reported measures,
1554 with objective indicators of behavior.

1555 In terms of research perspectives, future studies should in-
1556 vestigate intention–automaticity profiles in different age groups,
1557 because both the profiles and the patterns of behaviors at-
1558 tached to them could differ among adolescents (younger and
1559 older), adults (depending on their working and family status),
1560 and older people. Particular attention should be paid to different

cultures to examine whether the current results can be generalized. Finally, in a more applied perspective, the study of these different psychological and behavioral profiles offers at least 2 new perspectives. First, specific interventions targeting preferentially controlled or automatic processes were recently proposed.^{36,37} It would be interesting to examine whether the efficacy of these interventions regarding behavior varies depending on the participants' motivational profile. Second, there is increasing evidence relative to the importance of considering behavioral change interventions for PA and SB in an independent fashion.³⁸ Considering the distinct nature of these behaviors, it could be fruitful to examine whether the efficacy of an intervention targeting controlled or automatic processes differs for PA and SB.³⁹

5. Conclusion

In this set of studies four distinct profiles emerged regarding the combination of PA and screen-based SB motivation in French adolescents. This result highlights the fact that individuals may display more or less high levels of intentions and/or automaticity toward these behaviours. The data further indicate that their adoption is rather independent. The healthiest behavioural pattern was displayed by those with a PA motivational profile, characterized by high levels of motivation toward PA but low toward screen-based SB, whereas the poorest was demonstrated by adolescents with the Screen profile, showing reverse characteristics. Future research should seek to replicate these findings, taking into account the type of behaviours adopted—structured or spontaneous—using both self-reported and objective measurement, and exploring whether the results differ according to age or culture. In a translational perspective, the response of individuals displaying various motivational profiles to interventions targeting preferentially intentional or automatic motivation toward PA and/or both SB could be investigated.

Acknowledgement

The authors are grateful to Annick Spiekermann and Marco Neuvel for help in the data collection process.

Authors contribution

GM drafted the manuscript; GC performed the statistical analysis and drafted the manuscript; JB conceived of the study, participated in its design and coordination and helped drafting the manuscript. All authors have read and approved the final version of the manuscript, and agree with the order of presentation of the authors.

Competing interests

None of the authors declare competing financial interests.

References

- Hallal PC, Andersen LB, Bull FC, Guthold R, Haskell W, Ekelund U. Global physical activity levels: surveillance progress, pitfalls, and prospects. *Lancet* 2012;**380**:247–57.
- Dumith SC, Gigante DP, Domingues MR, Kohl HW 3rd. Physical activity change during adolescence: a systematic review and a pooled analysis. *Int J Epidemiol* 2011;**40**:685–98.
- Mutz DC, Roberts DF, Van Vuuren DP. Reconsidering the displacement hypothesis: television's influence on children's time use. *Commun Res* 1993;**20**:51–75.
- Väistö J, Eloranta AM, Viitasalo A, Tompuri T, Lintu N, Karjalainen P. Physical activity and sedentary behaviour in relation to cardiometabolic risk in children: cross-sectional findings from the Physical Activity and Nutrition in Children (PANIC) Study. *Int J Behav Nutr Phys Act* 2014;**11**:55. doi:10.1186/1479-5868-11-55
- Pearson N, Braithwaite RE, Biddle SJ, van Sluijs EM, Atkin AJ. Associations between sedentary behaviour and physical activity in children and adolescents: a meta-analysis. *Obes Rev* 2014;**15**:666–75.
- Schmitz KH, Lytle LA, Phillips GA, Murray DM, Birnbaum AS, Kubik MY. Psychosocial correlates of physical activity and sedentary leisure habits in young adolescents: the Teens Eating for Energy and Nutrition at School study. *Prev Med* 2002;**34**:266–78.
- Bargh JA. The four horsemen of automaticity: Awareness, efficiency, intention, and control in social cognition. In: Wyer RS Jr, Srull TK, editors. *Handbook of social cognition*. 2nd ed. Hillsdale, NJ: Erlbaum; 1994.p.1–40.
- Moors A, De Houwer J. Automaticity: a conceptual and theoretical analysis. *Psychol Bull* 2006;**132**:297–326.
- Hagger MS, Rebar AL, Mullan B, Lipp OV, Chatzisarantis NL. The subjective experience of habits captured by self-report indexes may lead to inaccuracies in the measurement of habitual action. *Health Psychol Rev* 2015;**9**:296–302.
- Hollands GJ, Marteau TM, Fletcher PC. Non-conscious processes in changing health-related behaviour: a conceptual analysis and framework. *Health Psychol Rev* 2016;**1**–14.
- Verplanken B, Melkevik O. Predicting habit: the case of physical exercise. *Psychol Sport Exerc* 2008;**9**:15–26.
- Plotnikoff RC, Costigan SA, Karunamuni N, Lubans DR. Social cognitive theories used to explain physical activity behavior in adolescents: a systematic review and meta-analysis. *Prev Med* 2013;**56**:245–53.
- Ajzen I. The theory of planned behavior. *Organ Behav Hum Decis Process* 1991;**50**:179–211.
- Hagger MS, Chatzisarantis N, Biddle SJH. A meta-analytic review of the theories of reasoned action and planned behavior in physical activity: predictive validity and the contribution of additional variables. *J Sport Exerc Psychol* 2002;**24**:3–32.
- McEachan RRC, Conner M, Taylor NJ, Lawton RJ. Prospective prediction of health related behaviours with the Theory of Planned Behaviour: a meta-analysis. *Health Psychol Rev* 2011;**5**:97–144.
- Rhodes RE, Mark RS, Temmel CP. Adult sedentary behavior: a systematic review. *Am J Prev Med* 2012;**42**:e3–28.
- Marteau TM, Hollands GJ, Fletcher PC. Changing human behavior to prevent disease: the importance of targeting automatic processes. *Science* 2012;**337**:1492–5.
- Verplanken B, Orbell S. Reflections on past behavior: a self-report index of habit. *J Appl Soc Psychol* 2003;**33**:1313–30.
- Gardner B, de Bruijn GJ, Lally P. A systematic review and meta-analysis of applications of the self-report habit index to nutrition and physical activity behaviours. *Ann Behav Med* 2011;**42**:174–87.
- de Bruijn GJ, Van den Putte B. Adolescent soft drink consumption, television viewing and habit strength. Investigating clustering effects in the Theory of Planned Behaviour. *Appetite* 2009;**53**:66–75.
- Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;**35**:1381–95.
- Guedes DP, Lopes CC, Guedes JERP. Reproducibility and validity of the International Physical Activity Questionnaire in adolescents. *Rev Bras Med Esporte* 2005;**11**:147–54.
- Rosenberg DE, Norman GJ, Wagner N, Patrick K, Calfas KJ, Sallis JF. Reliability and validity of the Sedentary Behavior Questionnaire (SBQ) for adults. *J Phys Act Health* 2010;**7**:697–705.
- Deforche B, van Dyck D, Deliens T, de Bourdeaudhuij I. Changes in weight, physical activity, sedentary behaviour and dietary intake during the transition to higher education: a prospective study. *Int J Behav Nutr Phys Act* 2015;**12**:16. doi:10.1186/s12966-015-0173-9

- 691 25. Gardner B, Abraham C, Lally P, de Bruijn GJ. Towards parsimony in habit
692 measurement: testing the convergent and predictive validity of an
693 automaticity subscale of the Self-Report Habit Index. *Int J Behav Nutr*
694 *Phys Act* 2012;**9**:102. doi:10.1186/1479-5868-9-102
- 695 26. Tabachnick BG, Fidell LS. *Using multivariate statistics*. Boston, MD:
696 Pearson; 2013.
- 697 27. Hair JF, Anderson RE, Tatham RL, Black WC. *Multivariate data analysis*.
698 Upper Saddle River: Prentice Hall; 1998.
- 699 28. Boiché J, Marchant G, Nicaise V, Bison A. Development of the Generic
700 Multifaceted Automaticity Scale (GMAS) and preliminary validation for
701 physical activity. *Psychol Sport Exerc* 2016;**25**:60–7.
- 702 29. Melkevik O, Torsheim T, Iannotti RJ, Wold B. Is spending time in
703 screen-based sedentary behaviors associated with less physical activity:
704 a cross national investigation. *Int J Behav Nutr Phys Act* 2010;**7**:46.
705 doi:10.1186/1479-5868-7-46
- 706 30. Verplanken B, Friberg O, Wang CE, Trafimow D, Woolf K. Mental habits:
707 metacognitive reflection on negative self-thinking. *J Pers Soc Psychol*
708 2007;**92**:526–41.
- 709 31. Gardner B. Defining and measuring the habit impulse: response to
710 commentaries. *Health Psychol Rev* 2015;**9**:318–22.
- 711 32. Hagger MS, Rebar AL, Mullan BM, Lipp OV, Chatzisarantis NLD. The
712 subjective experience of habit captured by self-report indexes may lead to
713 inaccuracies in the measurement of habitual action. *Health Psychol Rev*
714 2015;**9**:296–302.
33. Keatley DA, Chan DK, Caudwell K, Chatzisarantis NL, Hagger MS. A
715 consideration of what is meant by automaticity and better ways to measure
716 it. *Front Psychol* 2015;**5**:1537. doi:10.3389/fpsyg.2014.01537 717
34. Che S, Zhu X, Welk GJ, Kim Y, Lee J, Meier NF. Using Sensewear
718 armband and diet journal to promote adolescents' energy balance
719 knowledge and motivation. *J Sport Health Sci* 2014;**3**:326–32. 720
35. Perugini M, Richetin J, Zogmeister C. Prediction of behavior. In:
721 Gawronski B, Payne K, editors. *Implicit social cognition*. New York:
722 Guilford Press; 2010.p.255–77. 723
36. Williams SL, Michie S, Dale J, Stallard N, French DP. The effects of a brief
724 intervention to 428 promote walking on Theory of Planned Behavior
725 constructs: a cluster randomized controlled trial in 429 general practice.
726 *Patient Educ Couns* 2015;**98**:651–9. 727
37. Lally P, Gardner B. Promoting habit formation. *Health Psychol Rev*
728 2013;**7**:137–58. 729
38. Martin A, Fitzsimons C, Jepson R, Saunders DH, van der Ploeg HP,
730 Teixeira PJ, et al. Interventions with potential to reduce sedentary time in
731 adults: systematic review and meta-analysis. *Br J Sports Med* 2015;
732 **49**:1056–63. 733
39. Rhodes RE, Nigg CR. Advancing physical activity theory: a review and
734 future directions. *Exerc Sport Sci Rev* 2011;**39**:113–19. 735